



# **Technical Datasheet**

# **HiGlutaXL™ Minimum Essential Medium Eagle**

(Alpha modification)

With L-Alanyl-L-Glutamine, Sodium bicarbonate Without Deoxyribonucleosides and Ribonucleosides

**Product Code: AL081G** 

## **Product Description:**

 $HiGlutaXL^{TM}$ medium contains the stabilised dipeptide form of L-glutamine, L-alanyl-L-glutamine. HiGlutaXL<sup>TM</sup> medium offers several advantages over the conventional glutamine containing media. Dipeptide form prevents the intramolecular cyclization reaction, thus preventing toxic build-up of ammonia. L-alanyl-Lglutamine incorporates L-alanine that protects the alpha amino acid group. Aminopeptidases within the cell break the dipeptide, gradually releasing both L-glutamine and L-alanine for use by the cell. The gradual release of Lglutamine obviates the need to supplement L-glutamine frequently and allows liquid media to be stored at 4°C for longer periods.

AL081G is alpha modification of HiGlutaXL $^{\text{TM}}$  Minimum Essential Medium Eagle with L-alanyl-L-glutamine and sodium bicarbonate. It is devoid of deoxyribonucleosides and ribonucleosides. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

### **Composition:**

| Ingredients                           | mg/L     |
|---------------------------------------|----------|
| INORGANIC SALTS                       |          |
| Calcium chloride dihydrate            | 265.000  |
| Magnesium sulphate anhydrous          | 97.720   |
| Potassium chloride                    | 400.000  |
| Sodium bicarbonate                    | 2200.000 |
| Sodium chloride                       | 6800.000 |
| Sodium dihydrogen phosphate anhydrous | 122.000  |
| AMINO ACIDS                           |          |
| Glycine                               | 50.000   |
| L-Alanine                             | 25.000   |
| L-Alanyl-L-Glutamine                  | 434.000  |
| L-Arginine hydrochloride              | 126.000  |
| L-Asparagine monohydrate              | 44.000   |
| L-Aspartic acid                       | 30.000   |
| L-Cysteine hydrochloride monohydrate  | 100.000  |

| L-Cystine dihydrochloride             | 31.300   |
|---------------------------------------|----------|
| L-Glutamic acid                       | 75.000   |
| L-Histidine hydrochloride monohydrate | 42.000   |
| L-Isoleucine                          | 52.000   |
| L-Leucine                             | 52.000   |
| L-Lysine hydrochloride                | 72.500   |
| L-Methionine                          | 15.000   |
| L-Phenylalanine                       | 32.000   |
| L-Proline                             | 40.000   |
| L-Serine                              | 25.000   |
| L-Threonine                           | 48.000   |
| L-Tryptophan                          | 10.000   |
| L-Tyrosine disodium salt dihydrate    | 51.900   |
| L-Valine                              | 46.000   |
| VITAMINS                              |          |
| Choline chloride                      | 1.000    |
| D-Biotin                              | 0.100    |
| D-Ca-Pantothenate                     | 1.000    |
| Folic acid                            | 1.000    |
| L-Ascorbic acid                       | 50.000   |
| Nicotinamide                          | 1.000    |
| Pyridoxal hydrochloride               | 1.000    |
| Riboflavin                            | 0.100    |
| Thiamine hydrochloride                | 1.000    |
| Vitamin B12                           | 1.360    |
| i-Inositol                            | 2.000    |
| OTHERS                                |          |
| D-Glucose                             | 1000.000 |
| Lipoic acid                           | 0.200    |
| Phenol red sodium salt                | 11.000   |
| Sodium pyruvate                       | 110.000  |
|                                       |          |

### **Quality Control:**

**Appearance** 

Orangish red colored, clear solution.

pН

7.00 - 7.60

Osmolality in mOsm/Kg H<sub>2</sub>O

280.00 -320.00

#### **Sterility**

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

#### **Cultural Response**

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts.

#### **Endotoxin Content**

NMT 1EU/ml

### **Storage and Shelf Life:**

Store at 2-8°C away from bright light. Shelf life is 18 months. Use before expiry date given on the product label.

Disclaimer: Revision: 04/2022

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia<sup>TM</sup> publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia<sup>TM</sup> Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

