

## CELLin1™

**Chemically defined, Animal Component Free Virus Production medium**

**With Sodium bicarbonate**

**Without L-Glutamine**

**Contains basal medium in liquid form**

**Product Code: SFM036L**

### Product description:

SFM036L, CELLin1™ is chemically defined, animal component-free medium designed for growth and maintenance of Vero, MDCK, MDBK, PK15 and MRC-5 cells. The medium is devoid of L-Glutamine.

### Contents:

| Code   | Contents             |
|--------|----------------------|
| Part A | Basal Medium, liquid |
| Part B | Growth Supplement    |

### Directions:

#### Preparation of complete medium (SFM036L):

1. Thaw the growth supplement (Part B) overnight at 2-8°C.
2. Disinfect the external surface of bottle of Part A and Part B by spraying isopropyl alcohol before placing in a biosafety hood.
3. Transfer the entire content of one bottle of Part B to given quantity of basal medium (Part A) under aseptic conditions.

*Note: If desired, 10ml of Antibiotic-Antimycotic solution (A002) can be added to 1 litre of complete medium.*

4. Add 20ml/L of 200mM L-Glutamine solution (TCL012) to the medium to obtain final concentration of 4mM L-Glutamine.

*Note: Do not mix vigorously. Doing so will cause formation of foam.*

5. Store the complete medium (SFM036L) at 2-8°C until use.

#### Procedure for Adaptation:

Cells are adapted to serum free conditions by gradual weaning. Gradual weaning is a slow procedure that involves decreasing the percentage of serum in the medium, thereby gradually adapting the cells to serum free conditions.

### Critical points:

- Cells used for adaptation should exhibit a healthy morphology and have more than 90% viability.
- Cells should be in the mid-logarithmic phase of growth.
- It is necessary to subculture the cells at least thrice at each step, before going to the next step of adaptation.
- Subculturing should be performed when the cells are 70-80% confluent.

### Gradual weaning:

*Note: The procedure for gradual weaning is also applicable for adapting cells from current serum free medium to the new serum free medium.*

1. Subculture the cells from serum containing medium and seed them in 75:25 ratio of serum containing medium and SFM036L with a seeding density of 0.3-0.5X10<sup>6</sup> cells/ml.
2. Incubate at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>. Make provision for gas exchange by loosening the caps of flasks in case of closed caps or use vented caps.
3. Subculture once the cells are 70-80% confluent.
4. Determine cell density and reseed the cells in 75:25 ratio of serum containing medium and SFM036L.

*Note: It is necessary to subculture the cells at least thrice at each step of adaptation before going to the next step.*

5. Repeat steps 1 to 4 for three subcultures of each step of gradual adaptation.

*Note: Refer figure 1 for details of each adaptation step.*

6. After step 3 (25:75 serum containing medium: SFM036L) of adaptation, the cells cannot be directly subcultured in 100% serum free conditions. Complete withdrawal of serum may alter cell morphology and decrease the cells viability. Hence, it is very critical to maintain them at 10:90, 5:95 and 1:99 ratios before 100% serum free conditions

7. When the cells reach 100% serum free step of adaptation, subculture them respectively till a cell density of  $1.5- 2.0 \times 10^6$  cells/ml is obtained within 4-6 days of culture. At this point, the cells are considered to be adapted to serum free conditions.

### Material required but not provided:

Tissue culture grade water (TCL010)  
 L-Glutamine powder (TC243)  
 L-Glutamine 200mM Solution (TCL012)  
 Sodium bicarbonate powder (TC230)  
 7.5% Sodium bicarbonate solution (TCL013)  
 1N Hydrochloric acid (TCL003)  
 1N Sodium hydroxide (TCL002)  
 Trypsin – EDTA solution 1X (TCL007)  
 EnVzyme™ Easy (TCL137)  
 Trypsin Inhibitor from Soyabean 1X (TCL068)

### Quality control:

#### Appearance

Part A: Orangish red colored clear solution  
 Part B: Clear colorless to pale yellow solution

#### pH of Part A

7.00 - 7.60


#### Osmolality of Part A (mOsm/Kg H<sub>2</sub>O)

280.00- 320.00

#### Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium.

Fig 1: Gradual weaning



### Storage and shelf life:

Store basal medium at 2-8°C away from bright light.

Store serum free growth supplement at -20 °C.

Use before expiry date given on the product label.

Shelf life of the complete medium is 8 weeks at 2-8°C.

*Note: Freezing of the basal medium and complete medium is not recommended. Avoid repeated freezing and thawing of the growth supplement.*

Revision No: 0 / 2019

### Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ Publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.